

Earthquakes in the New Madrid Seismic Zone— What should we expect?

Randall W. Jibson

U.S. Geological Survey
Golden, Colorado




Most earthquakes occur along
plate boundaries, but not all
of them!



Seismicity of the US, 1977-1997



New Madrid Seismic Zone: Important Geologic Structures



widely felt

M-5.4 Alum Rock, CA, 30 Oct 2007


M-5.2 Illinois, 18 Apr 2008



INTENSITY	I	II-III	IV	V	VI	VII	VIII	IX	X+
SHAKING	Not felt	Weak	Light	Moderate	Strong	Very strong	Severe	Violent	Extreme
DAMAGE	none	none	none	Very light	Light	Moderate	Moderate/Heavy	Heavy	Very Heavy

INTENSITY	I	II-III	IV	V	VI	VII	VIII	IX	X+
SHAKING	Not felt	Weak	Light	Moderate	Strong	Very strong	Severe	Violent	Extreme
DAMAGE	none	none	none	Very light	Light	Moderate	Moderate/Heavy	Heavy	Very Heavy

1811 versus 1906




Area within Intensity VII:

New Madrid = 203,000
square miles

San Francisco = 12,000
square miles

The 1811-1812 New Madrid Earthquakes

Mainshocks and significant aftershocks




What happened in 1811-1812?

- Felt over 2 million square miles
 - ✓ Canada, East Coast, Florida
- Damage occurred over 230,000 square miles
- Geologic deformation—river-bank collapses, subsidence, uplift, liquefaction, and landslides—occurred over about 50,000 square miles
 - ✓ Cairo to Memphis, Crowleys Ridge to the Chickasaw bluffs


Geologic deformation

River-bank collapse



Geologic deformation

River-bank collapse




Geologic deformation

Subsidence



Geologic deformation


Uplift



Geologic deformation

Landsliding

- More than 220 large landslides mapped along about 200 miles of bluffs from Kentucky to Mississippi
- Other landslides were likely triggered up to 250 miles from the epicenters




Geologic deformation

Landsliding




Geologic deformation

Landsliding



Geologic deformation


Liquefaction



Geologic deformation

Liquefaction and sand blow formation:

- During earthquake, water-saturated sand is shaken.
- If shaking is strong and lasts long enough, pore-water pressure builds up; the sand loses its strength and acts like a liquid.
- A pressurized slurry of water and sand erupts to the surface, forming



Geologic deformation

liquefaction and
lateral spreading



Geologic deformation

Liquefaction



Some facts about the New Madrid Seismic Zone:

- New Madrid Seismic Zone produced large quakes in 1811-12, ~1450 AD, ~900 AD, and ~2350 BC
- During past 1200 years, the average time between these events is about 500 years
- The prehistoric earthquakes were similar in size to the 1811-1812 earthquakes
- Each New Madrid event was a sequence of earthquakes, including multiple very large mainshocks, much like the 1811-1812 sequence

50-year earthquake probabilities in the New Madrid Seismic Zone


- Repeat of 1811-1812 (M 7.5-8.0)
 - 7-10%
- Magnitude 6.0 or greater
 - 28-46%
- Magnitude 5.0 or greater
 - High probability

What will happen in a future
major earthquake?

“The past is the key to the
future”

Strong shaking

-- Earthquake Planning Scenario --
 ShakeMap for Nle2011nmsw7.7 Scenario
 Scenario Date: Mon May 16, 2011 14:00:00 GMT M 7.7 N35.90 W89.90 Depth: 10.0km




PLANNING SCENARIO ONLY -- Map Version 19 Processed Tue May 3, 2011 08:59:15 AM MDT

PERCEIVED SHAKING	Not felt	Weak	Light	Moderate	Strong	Very strong	Severe	Violent	Extreme
POTENTIAL DAMAGE	none	none	none	Very light	Light	Moderate	Moderate/Heavy	Heavy	Very Heavy
PEAK ACC. (%g)	<.17	.17-1.4	1.4-3.9	3.9-9.2	9.2-18	18-34	34-65	65-124	>124
PEAK VEL. (cm/s)	<0.1	0.1-1.1	1.1-3.4	3.4-8.1	8.1-16	16-31	31-60	60-116	>116
INSTRUMENTAL INTENSITY	I	II-III	IV	V	VI	VII	VIII	IX	X+

Strong shaking

-- Earthquake Planning Scenario --
Peak Accel. Map (in %g) for Nle2011nmsw7.7 Scenario
Scenario Date: Mon May 16, 2011 14:00:00 GMT M 7.7 N35.90 W89.90 Depth: 10.0km



PLANNING SCENARIO ONLY -- Map Version 19 Processed Tue May 3, 2011 08:59:15 AM MDT

Building damage





River-bank collapse

- Extensive bank failures for perhaps 100 miles in either direction from the epicenter
- Huge amounts of timber and debris in the river




Subsidence

- Thousands of square miles subside as much as 15 feet
- Flooding of subsided areas




Uplift




Landsliding

- Large landslides on bluffs from Cairo to Memphis and on Crowleys Ridge




Landsliding

- Large landslides on bluffs from Cairo to Memphis and on Crowley's Ridge
- Other landslides as far as 250 miles from the epicenter





Liquefaction

- More than 50,000 square miles could be affected
- As much as 6 feet of sand could be deposited on the



Effects

Agriculture



Effects

Roads



Effects

Roads



Effects

Bridges



Effects

Railroads



Effects

Rivers



Effects

Rivers



Effects


Levees



2/15/04 5:21
Earthquake damage
Looking Downstream, Salt

Effects

Ports



Effects

Power lines




Effects

Undergro
und
utilities



Effects

Fuel



Summary of the hazard

- The NMSZ experienced multiple magnitude-7+ earthquakes in 1811-1812 and at least three prior times in the last 5,000 years.
- The NMSZ is currently producing earthquakes; it is the most active zone in eastern North America.
- Large earthquake sequences have occurred about every 500 years in the recent past.
- A magnitude-7+ earthquake is a low-probability, high-consequence event; a magnitude-6+ earthquake has higher probability and also can cause significant damage.
- The consequences of a future major earthquake sequence in the NMSZ will be very serious if not catastrophic.

Sources of additional information

- Real-time and background information on earthquakes *earthquake.usgs.gov*
- New Madrid bicentennial web site:
newmadrid2011.org
- National Earthquake Hazards Reduction Program:
www.nehrp.gov



jibson@usgs.gov

